	-	FOR OFFICIAL						-
•			UJL					
			1					
		Nationa Qualific		nc				
		2021 AS			ESOI	JRCE	Mar	k
	X813/75/01						Che	mistry
				9	Sect	ion 1 –	– Answ	
	Duration — 2 hours 30 min	utes				*	X 8 1 3	7501*
	Fill in these boxes and read	l what is prii	nted b	elow.				
	Full name of centre				Town			
	Forename(s)	ç	urnam				Numbe	r of seat
								l of seat
	Date of birth Day Month	Voor		Scottish can	didate	numbor		
	Day Month	Year		Scottish can				
	Total marks — 100		_					
	SECTION 1 — 25 marks							
	Attempt ALL questions. Instructions for the completion	tion of Sectio	on 1 ar	e given on <i>i</i>	oage ()	2.		
	SECTION 2 — 75 marks			5 1	5			
	Attempt ALL questions.							
	You may refer to the Chemi Write your answers clearly and rough work is provided identify the question numb booklet. You should score t	n the spaces at the end o er you are at	s provi of this tempt	ded in this I booklet. If y ing. Any rou	bookle ou use igh wo	e this space rk must be	you must c written in t	learly his
	Use <b>blue</b> or <b>black</b> ink. Before leaving the examina Invigilator; if you do not, yo						×	SQA
	-		<b>0</b> 1 7					©

Downloaded free from https://sqa.my/

The questions for Section 1 are contained in the question paper X813/75/02.

Read these and record your answers on the answer grid on page 03 opposite.

Use **blue** or **black** ink. Do NOT use gel pens or pencil.

- 1. The answer to each question is **either** A, B, C or D. Decide what your answer is, then fill in the appropriate bubble (see sample question below).
- 2. There is **only one correct** answer to each question.
- 3. Any rough working should be done on the additional space for answers and rough work at the end of this booklet.

#### Sample question

To show that the ink in a ball-pen consists of a mixture of dyes, the method of separation would be


- A fractional distillation
- B chromatography
- C fractional crystallisation
- D filtration.

The correct answer is  $\mathbf{B}$  — chromatography. The answer  $\mathbf{B}$  bubble has been clearly filled in (see below).



#### Changing an answer

If you decide to change your answer, cancel your first answer by putting a cross through it (see below) and fill in the answer you want. The answer below has been changed to **D**.



If you then decide to change back to an answer you have already scored out, put a tick ( $\checkmark$ ) to the **right** of the answer you want, as shown below:









	Α	В	С	D
1	0	0	0	0
2	0	0	0	0
3	0	0	0	0
4	0	0	0	0
5	0	0	0	0
6	0	0	0	0
7	0	0	0	0
8	0	0	0	0
9	$\bigcirc$	$\bigcirc$	$\bigcirc$	0
10	0	0	0	0
11	0	$\bigcirc$	$\bigcirc$	0
12	0	0	0	0
13	0	0	0	0
14	0	0	0	0
15	0	0	0	0
16	0	0	0	0
17	0	0	0	0
18	0	0	0	0
19	0	0	0	0
20	0	0	0	0
21	0	0	0	0
22	0	0	0	0
23	0	$\bigcirc$	$\bigcirc$	0
24	0	0	0	0
25	0	0	0	0



[Turn over

Γ

L

DO NOT WRITE ON THIS PAGE



						MARKS	DO NO WRITE THIS	
	SEC	TION 2 — 75 m	arks				MARGI	
	Atte	empt ALL quest	ions					
The	e element tin has the che	mical symbol Sr	1.					
(a)	A sample of tin contains for each is shown.	three different	t isotopes.	The nuclic	le notation			
	<sup>116</sup> 50	<sup>118</sup> 50	n	<sup>120</sup> 50Sn				
	(i) State what is mea	nt by the term i	isotope.			1		
	(ii) This sample of tin State the mass nu sample of tin.	_				1		
(b)	Another isotope of tin e Write the nuclide notati					1		
(c)	Tin(IV) chloride can be formed by reacting tin with chlorine. Some properties of tin(IV) chloride are shown in the table.							
	Melting point		_33 ℃					
	Boiling point		114 °C					
	Electrical conductivity	as a solid	Does not	conduct				
	Electrical conductivity	as a liquid	Does not	conduct				
	Using the information in the table, state the <b>type of bonding</b> present in tin(IV) chloride.							
					[Turn ove	r		

Γ



2. A student carried out an investigation into reaction rates using dilute hydrochloric acid and indigestion tablets which contain calcium carbonate.

 $2HCl(aq) + CaCO_3(s) \rightarrow CaCl_2(aq) + H_2O(\ell) + CO_2(g)$ 

THIS


1

1


1

(a) Complete the diagram to show the apparatus required to collect and measure the volume of gas produced.

(An additional diagram, if required, can be found on page 29.)



(b) The student recorded the volume of gas produced over a period of time. A graph of the results of this experiment is shown.



(i) Add a curve to the graph to show the results that would be expected if the experiment was repeated using a crushed indigestion tablet.

(An additional diagram, if required, can be found on page 29.)

(ii) As these reactions proceed the rate of reaction decreases. Suggest a reason why the rate of reaction decreases.



### 2. (continued)

(c) The student carried out another three experiments, recording the time taken for 50 cm<sup>3</sup> of gas to be collected at different temperatures.

The results are shown.

Experiment	Temperature of acid (°C)	Time taken for 50 cm <sup>3</sup> of gas to be collected (s)
1	15	230
2	25	145
3	35	76

(i) Calculate the average rate of reaction, in  $\text{cm}^3 \text{s}^{-1}$ , for **experiment 1**. **2** 

(ii) State the relationship between temperature of acid and time taken to collect 50 cm<sup>3</sup> of gas.

(iii) **Experiment 1** was repeated using  $1 \cdot 0 \mod l^{-1}$  sulfuric acid,  $H_2SO_4(aq)$ , instead of  $1 \cdot 0 \mod l^{-1}$  hydrochloric acid, HCl(aq). The time taken to collect 50 cm<sup>3</sup> of gas decreased. Explain why the time taken decreased.



page 07

MARKS DO NOT WRITE IN THIS MARGIN

1

1

[Turn over

3.	Ami	monia	is a starting material for the commercial production of nitric acid.	MARKS	DO NOT WRITE IN THIS MARGIN
	(a)		alyst is used in the production of nitric acid. what is meant by the term catalyst.	1	
	(b)	Ammonia and nitric acid react together to form ammonium nitrate. Ammonium nitrate is commonly used as a fertiliser because it contains the element nitrogen, which is essential for healthy plant growth. (i) Name another element essential for healthy plant growth.		1	
		(ii)	Describe another property of ammonium nitrate that makes it suitable for use as a fertiliser. You may wish to use the data booklet to help you.	1	
	(c)		her common fertiliser is urea, (NH <sub>2</sub> ) <sub>2</sub> CO. Calculate the percentage by mass of nitrogen in urea, (NH <sub>2</sub> ) <sub>2</sub> CO. <b>Show your working clearly.</b>	3	
		(ii)	Urea dissolving in water is an endothermic process. Suggest a piece of apparatus that could be used to confirm this process is endothermic.	1	

L



[Turn over for next question

DO NOT WRITE ON THIS PAGE




4. Read the passage below and answer the questions that follow.

#### Air Fresheners

There are three ways an air freshener can remove an unpleasant smell. These are:

- Overpower it with a stronger smell
- Disguise it by mixing it with molecules to create a pleasant smell
- Absorb it

The following molecules are often found in unpleasant toilet smells.




3-methylbutanoic acid

skatole

Some other molecules that make up these bad smells can contain sulfur atoms. For example, hydrogen sulfide ( $H_2S$ ) is the gas associated with the smell of rotten eggs.

Air fresheners can contain molecules such as cyclodextrins that can absorb bad smells. Another molecule which is added for the same purpose is triethylene glycol.



triethylene glycol

Adapted from an article by John Emsley in Education in Chemistry, September 2007

(a) Cyclodextrin molecules absorb bad smells.

Name another molecule added to air fresheners to absorb bad smells.

1

THIS



•				MARKS	DO NOT WRITE IN THIS	
	4.	(cor	ntinued)		MARGIN	
		(b)	Draw a diagram, showing all outer electrons, of the molecule associated with the smell of rotten eggs.	1		
		(c)	Calculate the mass, in grams, of one mole of skatole.	1		
		(c)		·		
		(d)	Name the functional group circled on the triethylene glycol molecule.	1		
			[Turn over			
L						





MARKS DO NOT WRITE IN THIS MARGIN

1


## 4. (continued)

(e) Two branched carboxylic acid molecules are shown.





3-methylbutanoic acid




(i) Draw the structure of 2,4-dimethylpentanoic acid.




# 4. (e) (continued)

(ii) Carboxylic acids can be used to produce alkanes by a reaction that involves the loss of carbon dioxide.




3-methylbutanoic acid produces alkane X as shown.



3-methylbutanoic acid

Draw the structure of alkane X.

(f) The molecule shown is associated with the smell of wet dogs. It will decolourise bromine solution quickly.



State the term used to describe molecules that decolourise bromine solution quickly.


[Turn over

1




page 13

MARKS DO NOT WRITE IN THIS MARGIN








### 6. (continued)

(b) Co-polymers are polymers made using more than one type of monomer. Poly(ethylene-vinyl acetate) is a co-polymer used to make shower curtains and football studs. The monomers used to make it are shown.

MARKS DO NOT WRITE IN THIS MARGIN

1

1



(i) These monomer units join together by addition polymerisation.

State why these monomers can take part in addition polymerisation.

(ii) Draw the repeating unit formed when one molecule of **monomer A** joins with one molecule of **monomer B**.



- 7. When an acid and a base react together, water and a salt are formed.
  - (a) Acids and bases can be classified as strong or weak.

The salts formed, if soluble, will have a pH that depends on the strength of the acid and base used.

strong acid	+	strong base	$\rightarrow$	neutral salt	+	water
strong acid	+	weak base	$\rightarrow$	acidic salt	+	water
weak acid	+	strong base	$\rightarrow$	alkaline salt	+	water

Examples of strong and weak acids and bases are shown in the tables.

Ac	ids	Bases			
Strong acid	Weak acid		Strong base	Weak base	
hydrochloric acid	methanoic acid		sodium hydroxide	ammonium hydroxide	

(i) Methanoic acid reacts with sodium hydroxide.Name the salt formed.

(ii) Predict the pH of the salt solution formed when hydrochloric acid reacts with ammonium hydroxide.

1

1

MARKS DO NOT WRITE IN THIS MARGIN

[Turn over




7.	(	ntinue		MARKS	DO NOT WRITE IN THIS MARGIN
7.	(b)		The volume of an acid required to neutralise an accurately		
			measured volume of a base can be measured as follows.		
			1. Pipette 10 cm <sup>3</sup> of a base into a conical flask		
			2. Add 3 drops of indicator solution		
			<ol> <li>Add 0·1 mol l<sup>-1</sup> of an acid from a burette until the indicator changes colour</li> </ol>		
			State the name of this technique.	1	
		(ii)	To determine the concentration of a base, the titre volumes must be concordant.		
			State what is meant by the term concordant.	1	
	(c)	Salts	have a wide variety of uses.		
		(i)	The salt strontium chloride is used in fireworks.		
			State the colour of the flame that would be seen when a firework containing the salt strontium chloride is burned. You may wish to use the data booklet to help you.	1	
		(ii)	Another salt, barium sulfate, is used in some medical procedures. Write the formula, showing the charge on each ion, for barium sulfate.	1	


Γ



produced.1(ii) Propane and butane are members of the alkane homologous series.1State what is meant by the term homologous series.1(iii) Balance the equation for the combustion of butane.1 $C_4H_{10}$ + $O_2 \rightarrow CO_2$ + $H_2O$	p. 00		gas contains a mixture of the hydrocarbons prop pane and butane are burned, carbon dioxide an		MARKS
State what is meant by the term homologous series. (iii) Balance the equation for the combustion of butane. $C_4H_{10} + O_2 \rightarrow CO_2 + H_2O$ (b) During a camping trip a can of baked beans was heated by burning camping gas. $\frac{\text{Specific heat capacity of baked beans}}{\text{Specific heat capacity of baked beans}} = \frac{3 \cdot 6 \text{ kJ kg}^{-1} \circ \text{C}^{-1}}{16 \text{ Energy absorbed by the baked beans}} = \frac{3 \cdot 6 \text{ kJ kg}^{-1} \circ \text{C}^{-1}}{16 \text{ Comparature of baked beans}} = \frac{3 \cdot 6 \text{ kJ kg}^{-1} \circ \text{C}^{-1}}{17 \circ \text{C}}}$ Calculate the final temperature, in °C, of the baked beans using the	(a)	(i)		n dioxide has been	1
$C_4H_{10} + O_2 \rightarrow CO_2 + H_2O$ (b) During a camping trip a can of baked beans was heated by burning camping gas. $\frac{\text{Specific heat capacity of baked beans}}{\text{Specific heat capacity of baked beans}} \frac{3.6 \text{ kJ kg}^{-1} \text{ oC}^{-1}}{16 \text{ Energy absorbed by the baked beans}} \frac{76.32 \text{ kJ}}{76.32 \text{ kJ}}$ Temperature of baked beans before being heated 17 °C Mass of baked beans} 400 g Calculate the final temperature, in °C, of the baked beans using the		(ii)	•	-	
(b) During a camping trip a can of baked beans was heated by burning camping gas. Specific heat capacity of baked beans 3.6 kJ kg <sup>-1</sup> °C <sup>-1</sup> Energy absorbed by the baked beans 76.32 kJ Temperature of baked beans before being heated 17 °C Mass of baked beans 400 g Calculate the final temperature, in °C, of the baked beans using the		(iii)			1
Energy absorbed by the baked beans76.32 kJTemperature of baked beans before being heated17 °CMass of baked beans400 gCalculate the final temperature, in °C, of the baked beans using the	(b)	camp	ping gas.	ed by burning	
Temperature of baked beans before being heated17 °CMass of baked beans400 gCalculate the final temperature, in °C, of the baked beans using the					
Mass of baked beans400 gCalculate the final temperature, in °C, of the baked beans using the		-			
Calculate the <b>final</b> temperature, in °C, of the baked beans using the		Ene	rgy absorbed by the baked beans	76·32 kJ	
		Ene Terr	rgy absorbed by the baked beans perature of baked beans before being heated	76·32 kJ 17 ℃	
		Ene Terr	rgy absorbed by the baked beans perature of baked beans before being heated	76·32 kJ 17 ℃	







## 9. (b) (continued)

(iii) State why ionic compounds, like aluminium oxide, conduct electricity when molten.

(iv) During electrolysis, the following reactions take place.

MARKS DO NOT WRITE IN THIS MARGIN

1

1

Write the redox equation for the overall reaction.



кеа	d the passage and answer the questions that follow.	7
	Tungsten carbide	
18	ungsten has the chemical symbol W. It can be traced back to the <sup>3<sup>th</sup> century when it was first extracted from the ore wolframite. Tungsten as a very high melting point of 3422 °C.</sup>	
ch he	ungsten carbide, a compound of tungsten, was accidentally made by nemist Henri Moissan in 1896. In an attempt to make artificial diamond, e heated sugar and tungsten(III) oxide in a furnace. The sugar reacted ith the tungsten oxide to produce liquid tungsten carbide.	
	ungsten carbide has a melting point of 2870 °C and a boiling point of 000 °C and is three and a half times as dense as titanium.	
ht	Adapted from tps://eic.rsc.org/magnificent-molecules/tungsten-carbide/3008556.article	
(a)	State the name of the ore from which tungsten was first extracted.	1
(b)	Write the formula for the compound that was heated with sugar, in a furnace, to produce tungsten carbide.	1
(c)	Suggest a temperature, in °C, that Henri Moissan's furnace could have been operating at when tungsten carbide was accidentally made.	1
(d)	Calculate the density of tungsten carbide, in $g \text{ cm}^{-3}$ . You may wish to use the data booklet to help you.	2

[Turn over




- 11. Carbon-14 is an isotope of carbon that can be used to determine the age of materials.
  - (a) When a neutron is absorbed by a nitrogen-14 nucleus, a carbon-14 isotope is produced along with one other particle, **X**.

An equation for this is shown

$${}^{14}_{7}N + {}^{1}_{0}n \rightarrow {}^{14}_{6}C + X$$

Name particle X.

(b) The graph shows how the percentage of carbon-14 in a sample changes over a period of time.



(i) Use the graph to calculate the half-life, in years, of carbon-14.


1

1

THIS



11. (b)	(con	tinund)	MARKS	DO NOT WRITE IN THIS MARGIN
11. (b)		Use your answer to part (b) (i) to calculate the age, in years, of a bone found to contain $\frac{1}{16}$ of the original carbon-14 content.	2	
	(iii)	Another bone, believed to be over 100 000 years old, cannot be dated using levels of carbon-14.	4	
		Suggest why carbon-14 is unsuitable for dating this bone.	1	
		[Turn over		
L		* X 8 1 3 7 5 0 1 2 5 *		





12.	(CO)	itinued)					MARKS	THIS
	(c)	Another method for making	ng cycloa	lkanes is shown.				MARGIN
		C <sub>5</sub> H <sub>10</sub> Br <sub>2</sub> + 1,5-dibromopentane			+ so	2NaBr dium bromide		
		Calculate the mass, in gra cyclopentane.	ams, of so	odium required to p	produc	e 175 g of	3	
								1
	(d)	Cycloalkanes can experier		strain within their r	rings. 1	The ring strain		
	(d)	of some cycloalkanes is sh	nown.			The ring strain		
	(d)	of some cycloalkanes is sh	hown. kane	Total ring strain		The ring strain		
	(d)	of some cycloalkanes is sh	hown. kane pane	Total ring strain 132		The ring strain		
	(d)	of some cycloalkanes is sh Cycloalk cyclopro cyclopen	kane pane ntane	Total ring strain 132 25		The ring strain		
	(d)	of some cycloalkanes is sh	kane pane ntane	Total ring strain 132		The ring strain		
	(d)	of some cycloalkanes is sh Cycloalk cyclopro cyclopen	kane pane ntane otane	Total ring strain 132 25	(kJ)			
	(d)	of some cycloalkanes is sh Cycloalk cyclopro cyclopen cyclohep	hown. kane ppane ntane otane n =	Total ring strain 132 25 28 total ring s nber of carbons in	(kJ) train the cy	cloalkane	1	
	(d)	of some cycloalkanes is sh Cycloalk cyclopro cyclopen cyclohep Ring strain per carbo	hown. kane ppane ntane otane n =	Total ring strain 132 25 28 total ring s nber of carbons in	(kJ) train the cy	cloalkane		
	(d)	of some cycloalkanes is sh Cycloalk cyclopro cyclopen cyclohep Ring strain per carbo	hown. kane ppane ntane otane n =	Total ring strain 132 25 28 total ring s nber of carbons in	(kJ) train the cy	cloalkane		
	(d)	of some cycloalkanes is sh Cycloalk cyclopro cyclopen cyclohep Ring strain per carbo	hown. kane ppane ntane otane n =	Total ring strain 132 25 28 total ring s nber of carbons in	(kJ) train the cy	cloalkane	1	




- MARKS DO NOT WRITE IN THIS MARGIN
- **13.** Vinegar is a solution of ethanoic acid in water. Different types of vinegar can contain different concentrations of ethanoic acid.

Using your knowledge of chemistry, suggest how a student could determine which type of vinegar had the highest concentration of ethanoic acid.

3


## [END OF QUESTION PAPER]





#### ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK


Additional diagram for question 6 (a) (ii)






## ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK

MARKS DO NOT WRITE IN THIS MARGIN







#### MARKS DO NOT WRITE IN THIS MARGIN

## ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK



#### MARKS DO NOT WRITE IN THIS MARGIN

## ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK



Г

L

DO NOT WRITE ON THIS PAGE



DO NOT WRITE ON THIS PAGE



## DO NOT WRITE ON THIS PAGE

Acknowledgement of copyright

Question 4	Article is adapted from "Air fresheners" by John Emsley, from <i>Education in Chemistry</i> , 1st September 2007. Reproduced by kind permission of Dr John Emsley.			
	SQA has made every effort to trace the owners of copyright of this item and seek permissions. We are happy to discuss permission requirements and incorporate any missing acknowledgement. Please contact question.papers@sqa.org.uk.			
Question 10	Article is adapted from "Tungsten Carbine" by Matthew Gunther, from <i>Education in Chemistry</i> , 25th January 2018.			
	SQA has made every effort to trace the owners of copyright of this item and seek permissions. We are happy to discuss permission requirements and incorporate any missing acknowledgement. Please contact question.papers@sqa.org.uk.			

